

Patient Information

Chlamydial Infection in Females

What is chlamydial infection?

Chlamydia is one of the most common sexually transmitted diseases in the US. It can affect several parts of the body. In women the infections often occur in the urinary tract and female organs.

How does it occur?

The organisms that cause the infection are called Chlamydia trachomatis. They are similar to bacteria. The infection is usually passed from person to person during sexual intercourse. It can also be passed by other intimate contact with the genital or rectal area.

An infection can be passed from a pregnant woman to her baby during birth. Chlamydia can cause the baby to have eye infections or pneumonia.

What are the symptoms?

Possible symptoms include:

- abnormal vaginal discharge
- mild pain or discomfort when you urinate
- menstrual periods that are heavier than usual
- more painful periods
- abdominal pain
- spotting between periods or after sex.

Often there are no symptoms, especially early in the infection.

How is it diagnosed?

Your healthcare provider will ask about your symptoms. You will usually have a pelvic exam to get a sample from your cervix for testing. Sometimes a sample of urine can be tested instead.

Because chlamydia is one of the most common sexually

Patient Information

Chlamydial Infection in Females

transmitted diseases, healthcare providers recommend yearly tests for chlamydia in sexually active teens and young women up to age 25. A test for chlamydia is also recommended for any woman who has a new sex partner or multiple sexual partners.

How is it treated?

Taking antibiotics usually cures the infection. You may need to take more than one antibiotic, especially if there is a chance you have other infections, such as gonorrhea. Your sexual partner or partners should also have treatment even if they have no symptoms.

Make sure to tell your healthcare provider if you are or think you are pregnant, so he or she will prescribe the correct antibiotic.

How long will the effects last?

Without treatment the disease can cause serious problems, such as scarring of the fallopian tubes, tubal pregnancy, and infertility. With antibiotic treatment, the symptoms usually go away within a few days after you start taking the medicine.

How can I take care of myself?

- ☐ Follow the full treatment prescribed by your healthcare provider. This includes taking your medicine for as long as it is prescribed, even if your symptoms are gone before you have finished taking it.
- ☐ Take aspirin or acetaminophen to reduce pain.
- ☐ If you are taking the antibiotic tetracycline or doxycycline, avoid milk products 1 to 2 hours before and after you take the medicine. Also, avoid sun exposure. The medicine may cause you to be very sensitive to the sun and get a severe sunburn.

Patient Information

Chlamydial Infection in Females

- Follow your healthcare provider's instructions for follow-up visits and tests. Your provider will need to check that the infection is gone.
- Tell everyone with whom you have had sex in the last 3 months about your infection. They must also be treated, even if they have no symptoms. Do not have sex before both you and your partner have finished all the medicine and your provider says it is OK.
- Have a checkup every year. See your provider between checkups if you are having symptoms of vaginal infection or discomfort during sex.
- Call your healthcare provider right away if:
 - You develop severe abdominal pain.
 - You vomit and cannot keep the medicine down.
 - You develop a fever over 100°F (37.8°C).
 - You feel you are getting sicker rather than better.

How can I help prevent chlamydial infection?

- Reduce the risk of infection by always using latex or polyurethane condoms during foreplay and vaginal, oral, or anal intercourse.
- Have just 1 sexual partner who is not sexually active with anyone else. Make sure your partner has been tested for chlamydia and other sexually transmitted diseases.
- Even if you don't have symptoms but have had unprotected sex (without a condom), see your healthcare provider to be checked for sexually transmitted diseases. If you have been sexually assaulted and are at risk for having been infected with chlamydia, you should be treated to prevent infection.
- If you are pregnant, tell your healthcare provider so you can help prevent infection in the baby.

[Related Topics]

Chlamydial Infection in Males
Prevention of Sexually Transmitted Diseases

Patient Information

Chlamydial Infection in Females

Pregnancy Prevention for Young Men and Women

Developed by David W. Kaplan, MD, and Phyllis Cooper RN, MN.

Published by RelayHealth.

© 2008 RelayHealth and/or one of its affiliates. All Rights Reserved.

This content is reviewed periodically and is subject to change as new health information becomes available. The information is intended to inform and educate and is not a replacement for medical evaluation, advice, diagnosis or treatment by a healthcare professional.